

Wintering habitats of Eleonora's Falcons *Falco eleonora* in Madagascar

UGO MELLONE^{1*}, PASCUAL LÓPEZ-LÓPEZ^{1,2}, RUBEN LIMIÑANA^{1,3} and VICENTE URIOS¹

¹Estación Biológica Terra Natura, Grupo de Investigación Zoología de Vertebrados, Instituto Universitario de Investigación CIBIO, University of Alicante, Apdo 99, E-03080, Alicante, Spain, ²Cavanilles Institute of Biodiversity and Evolutionary Biology, Terrestrial Vertebrates Group, University of Valencia, C/Catedrático José Beltrán 2, E-46980 Paterna, Valencia, Spain and ³Instituto de Investigación en Recursos Cinegéticos (IREC), CSIC-UCLM-JCCM, Ronda de Toledo, s/n, E-13005 Ciudad Real, Spain

Capsule Eleonora's Falcons wintering in Madagascar selected degraded humid forests and cultivated areas close to pristine humid forest.

Aims To identify the habitat preferences of Eleonora's Falcon *Falco eleonora* on their wintering grounds in Madagascar, and to use this information to gain insights into the conservation priorities of this species.

Methods A total of 11 Eleonora's Falcons were captured in Spain in 2007 and 2008 and equipped with solar-powered satellite transmitters. We obtained information on five complete wintering events for three birds, two of them tracked for two consecutive years. Data were analyzed using geographic information system-based cartography.

Results The analyses showed a preference for degraded humid forests and cultivated lands within areas where pristine humid forests were the most abundant habitat type.

Conclusions Eleonora's Falcons could be taking advantage from a spill-over edge effect of their insect prey into cultivated and more open areas close to humid forests. However, the importance of humid forests for Eleonora's Falcons seems to be high. The current loss of this habitat in Madagascar is a cause for concern with respect to the conservation of this long-distance migratory falcon species.

A consistent body of recent literature suggests that, in contrast to resident bird species, populations of migratory species can be affected by events occurring in different parts of the world (Newton 2004). Moreover, it has recently been shown that populations of long-distance migratory birds are declining at a faster rate than short-distance migrants or resident species (Sanderson *et al.* 2006). Therefore, understanding the interactions between different periods of the annual cycle in migratory birds becomes of the greatest importance for their conservation (Martin *et al.* 2007, Bowlin *et al.* 2010).

Eleonora's Falcons *Falco eleonora* are long-distance migratory raptors, which breed mainly in islands of the Mediterranean Sea and winter mainly in Madagascar (Walter 1979). The species is listed in the Annex I of the Directive 2009/147/EC and constitutes a priority species for conservation (BirdLife International 2010). Unfortunately, long-term series of comparable

demographic data are lacking, but on the basis of recent surveys, the species seems to be stable or steadily increasing across its whole distribution range (Del Moral 2008, Dimalexis *et al.* 2008). Although the migratory routes have recently been mapped in detail (Gschweng *et al.* 2008, López-López *et al.* 2009, López-López *et al.* 2010, Mellone, López-López *et al.* 2011), little is known about the species' habitat requirements on its wintering grounds. Such information is key for the conservation of Eleonora's Falcons, since actions for habitat preservation, especially at wintering grounds, remain largely unimplemented (BirdLife International 2010).

Our knowledge of the ecology of migratory birds has been constrained by the difficulties in tracking individuals across seasons. Several methods such as traditional ringing and banding have been employed in recent decades (Berthold 2001). Nevertheless, advances in the miniaturization of satellite tracking devices are now making it possible to determine the connections between the different seasonal grounds of birds in a

*Correspondence author. Email: ugomellone@libero.it

way unimaginable few years ago (Wikelski *et al.* 2007). Other methods, such as genetic markers, stable isotopes (Hobson 2008) and geolocators (Rodríguez *et al.* 2009), although complementary, do not allow fine-tuned habitat selection analyses that are possible with satellite tracking. Nevertheless, wintering-focused analyses performed with this technology have received little attention in comparison with the greater body of research dealing with migratory routes, and hence very few studies have been published so far (Gerkmann & Meyburg 2009, Jiguet *et al.* 2011).

The island of Madagascar has been identified as one of the most important biodiversity hotspots in the world, which is strongly endangered owing to several threats – mainly habitat loss, including deforestation (Myers *et al.* 2000, Harper *et al.* 2007, Irwin *et al.* 2010). In this context, the identification and conservation of optimal habitats for long-distance migratory species that winter in Madagascar is of the utmost importance. Besides benefits for the species itself, it could be useful also for wider conservation aims, given that raptors prove to be, in certain conditions, good biological indicators (Sergio *et al.* 2005, 2008). With the aim of identifying habitats where conservation actions should be focused, we analyze here the habitat preferences of Eleonora's Falcons wintering in Madagascar. According to the International Action Plan for the species, the location of the wintering areas and the description of their ecological characteristics are urgently needed and have been considered an 'essential' priority (BirdLife International 1999, 2010).

METHODS

Study species

Eleonora's Falcons are colonial raptors, which breed mainly on Mediterranean islands, from Spain to Greece, including several islands located in the central Mediterranean (Italy), and the northern coast of Africa (Algeria and Libya) (Walter 1979). They also breed in a few colonies located in the Atlantic Ocean, including the Canary Islands (Spain) and Mogador (Morocco). The bulk of the population (80–90%) breeds in Greece (Dimalexis *et al.* 2008). Winter records are almost exclusively restricted to Madagascar (Walter 1979, Gschweng *et al.* 2008, Kassara *et al.* 2011), although some observations (mainly juveniles) have been recorded on the eastern coast of Africa (Kenya, Tanzania and Mozambique) (Gschweng *et al.* 2008, authors unpubl. data). Since individuals belonging

to different populations concentrate in such a narrow winter range, a comprehensive mapping and a complete description of habitat requirements at wintering areas is crucial in order to gain insight into conservation priorities for the species as a whole. In fact, to the best of our knowledge, only sporadic visual observations of Eleonora's Falcons in Madagascar have been reported so far (Thorstrom & René de Roland 2000, Zefania 2001). In this context, information based even on a small sample size of tracked birds could be useful, at least as a starting point for further investigations, which could be also applied to fieldwork planning.

Animal tagging and transmitter terminal programming

A total of 11 Eleonora's Falcons were trapped in Balearic (39°30'N 3°00'E) and Columbretes Islands (39°55'N 0°40'E), located in the western Mediterranean (Spain) (Fig. 1). Birds were trapped in autumn 2007 and 2008, and were equipped with Microwave Telemetry Inc. 9.5-g solar-powered transmitter terminals (López-López *et al.* 2009, López-López *et al.* 2010). During the wintering season, transmitters were programmed to collect data (coordinates, date and time) on a duty cycle of 12 hours on/58 hours off. Since the transmitters provide data of different accuracy, only high-quality locations (location classes 1–3; maximum error < 1 km) were used for the analyses (Soutullo *et al.* 2007).

Here, we only used wintering data from those individuals for which we recorded the complete wintering season (between arrival on the wintering grounds and the onset of the following pre-breeding migration). We used data belonging to three adult individuals, two of them tracked for two consecutive years (i.e. five different wintering events). Locations obtained less than one hour after the previous one were excluded from the analyses to avoid spatial autocorrelation and, when more than one location was available within an hour, we used the one of highest quality (Limiñana *et al.* 2008). On average, each wintering event accounted for 235 locations (range: 187–286, $n = 5$) spanning from November to April.

Habitat preferences and kernel estimation

To map the wintering areas in detail, we estimated the individuals' home-range core area of every wintering event through fixed kernel density contours (*sensu* Worton 1989), using the Animal Movement Extension




Figure 1. (a–e) Wintering locations (●) of three Eleonora's Falcons tracked by satellite-telemetry in northern Madagascar. Range sizes (km²) according to kernel density contours for five wintering events are as follows: (a) individual #80399, wintering event 2008/09; (b) #80399, wintering event 2009/10; (c) #80402, wintering event 2008/09; (d) #80402, wintering event 2009/10; (e) #80400, wintering event 2008/09. Dark green area represents humid forest, while pale green area represents degraded humid forest (following Moat and Smith [2007]). (f) Location of Eleonora's Falcon's tagging sites in the western Mediterranean and the wintering areas in Madagascar.

for ArcView, version 3.2 (Hooge & Eichenlaub 2000). We calculated the 25%, 50%, 75% and 95% fixed kernels using the least-squares cross-validation (LSCV) procedure (Silverman 1986). We chose the 50% kernel to represent the core areas after a detailed exploratory analysis, since it represents a reasonable trade-off between including a sufficient number of locations excluding outliers and, moreover, it allows comparisons with similar studies (Mellone, Yáñez *et al.* 2011). In our case, nearly 60% of overall locations were included within the core areas (range: 42–93%; Fig. 1). To assess individual fidelity to the wintering grounds, we calculated the inter-year overlap of the core areas of the two birds tracked during consecutive years. In order to describe habitat composition of wintering areas, we calculated the percentage of every habitat type within the core areas for each one of the five wintering events separately, using the geographic information system layers produced by the Madagascar

Vegetation Mapping Project (<http://www.vegmad.org>). These layers are based on the combination of remote sensing technology and field data collected by expert botanists (2003–06), and are the most accurate vegetation maps of Madagascar that are readily available (Moat & Smith 2007). Despite the ongoing changes in land cover in the country, it is unlikely that it could alter our results, owing to the short time-lapse between map compilation and the collection of our falcon tracking data. In fact, the deforestation rate of Madagascar forests has decreased from the beginning of the 1990s, in comparison with previous decades (Harper *et al.* 2007). We performed a habitat selection analysis to check whether Eleonora's Falcons occurred more frequently in certain habitats than expected by chance. To this end, we calculated a minimum convex polygon (MCP) for each wintering event (thus, three for the first year and two for the second one), which represents the maximum extension of available habitat, from which

birds actively select the preferred habitats (Worton 1995). We then generated 1000 randomly distributed points within each one of the five MCPs using 'Hawth's analysis tools' for ArcMap 9.2 (Beyer 2004), and assigned a habitat class to every random point, as well to as every real recorded location. To determine whether a bird preferred any particular habitat we used Monte Carlo tests (Manly 1997) to compare the observed frequency of real locations in different habitat types with the expected ones (Cadahía *et al.* 2010). The expected frequencies were obtained by sampling 1000 times, the same number of real bird locations from the list of random locations (Soutullo *et al.* 2008). Each individual wintering event was analyzed separately to avoid statistical pseudoreplication and no analysis including all the five wintering events at the same time was conducted. In addition, the analyses were first carried out for diurnal and nocturnal locations separately. Since the results did not differ between diurnal and nocturnal locations, we then pooled day and night data for further analyses, and here we present the results based on the complete data set. Monte Carlo analyses were conducted using Excel's PopTools 3.1 (Hood 2010). Pair-wise comparisons were two-tailed and the critical alpha level was set at $P < 0.05$.

RESULTS

Eleonora's Falcons' wintering areas were located in northern Madagascar (Fig. 1), where birds spent about five months of the year. Taking into account the core areas (50% fixed kernel contours), birds used one or two areas each wintering season (Fig. 1), with a median overall area of 606 km² (range: 246–6891 km², $n = 5$; Table 1). The two birds tracked during two consecutive wintering seasons used areas in their second year that they had used in the previous year, indicating site-fidelity. Bird #80399 used a smaller area during the second year that was completely within the one used in the first year (Fig. 1). On the other hand, bird #80402 used areas of similar size in both years (Table 1), with an overlap of 78.4% (Fig. 1).

Humid forest was the most abundant habitat within the core areas (pooling all data: 52.4% of the total surface area; Table 1). Habitat selection analyses showed that within individuals' MCPs, Eleonora's Falcons actively selected some habitat types, showing preferences for degraded humid forests and cultivations ($P < 0.005$ in the five cases; Table 1) and avoiding grasslands and humid forest ($P < 0.008$ in the five cases; Table 1). Furthermore, a detailed inspection

of maps showed that birds selected degraded humid forests and cultivated areas close to pristine humid forests (Fig. 1).

DISCUSSION

Wintering habitats of Eleonora's Falcons

Our results show that Eleonora's Falcons wintered in northern Madagascar, and were mostly located on the humid eastern slopes of the island, rather than on the dry western ones (Moat & Smith 2007) (Fig. 1). The eastern region of Madagascar is amongst the highest areas of the country, which accounts for the high rainfall levels that occur there. In tropical rainforests, higher rainfall levels may result in high numbers of insects (Wolda 1978) and consequently, falcons may select these sites rather than dry lower areas of the island owing to higher insect prey availability.

Eleonora's Falcons showed clear winter site-fidelity in consecutive years. Although this finding is based on a small sample size (two birds in only two consecutive years), our results are in agreement with previous reports concerning other raptor species during the wintering season (Alerstam *et al.* 2006, Strandberg *et al.* 2008, García-Ripollés *et al.* 2010, Limiñana *et al.* 2011), which suggest that falcons select areas at least in part on the basis of knowledge acquired in previous years. In addition, all three individuals used approximately the same area during the first wintering season (Fig. 1), which might suggest some degree of bird association during winter. This is in agreement with Zefania (2001), who reported Eleonora's Falcons associating in groups of 3–21 in Madagascar. These results, together with those of individuals tracked from other populations (Sardinia and Greece) which spent the winter in northern Madagascar (Gschweng *et al.* 2008, Kassara *et al.* 2011), suggests that our findings are consistent across years and might be applied to a larger proportion of Eleonora's Falcons during the wintering season in Madagascar. These results also suggest that the strength of migratory linkages (i.e. migratory connectivity) between breeding ranges and wintering grounds among different European populations of Eleonora's Falcon should be the subject of future research.

Within the core wintering areas roughly half of the habitat was pristine humid forest. However, results of habitat selection analyses showed an active preference by Eleonora's Falcons for degraded humid forest and cultivated areas (Table 1).

Table 1. Extension of the distinct habitats (km²) encompassed within the 50% kernel contours and significance levels of the habitat selection analysis (Monte Carlo tests) of three Eleonora's Falcons during five wintering events in Madagascar.

Individual ID	80399						80400			80402						Total	
	2008/09			2009/10			2008/09			2008/09			2009/10				
	Habitat	Area	%	P	Area	%	P	Area	%	P	Area	%	P	Area	%	P	Area
Water	1	0.01	x	0	0.00	x	0	0.00	x	0	0.00	ns	0	0.00	ns	1	0.01
Bare soil/rock	0	0.00	x	0	0.00	x	2	0.26	0.400	0	0.00	0.001	0	0.00	ns	0	0.00
Cultivation	1099	15.95	0.001 (+)	223	9.35	0.001 (+)	114	18.72	0.001 (+)	22	8.34	0.001 (+)	18	7.21	0.005 (+)	1362	13.92
Western dry forest	0	0.00	0.001	0	0.00	ns	8	1.33	0.007	0	0.00	ns	0	0.00	ns	0	0.00
Plateau grassland – wooded mosaic	152	2.21	0.001 (-)	40	1.70	0.001 (-)	65	10.78	0.001 (-)	1	0.48	0.001 (-)	1	0.34	0.001 (-)	195	1.99
Wooded grassland – bushland	708	10.28	0.001 (-)	377	15.81	0.006 (-)	73	12.05	0.001 (-)	10	3.67	0.001 (-)	8	3.22	0.002 (-)	1102	11.27
Wetlands	26	0.38	ns	14	0.59	0.017	0	0.00	x	0	0.00	ns	0	0.00	0.001	40	0.41
Humid forest	3538	51.34	0.001 (-)	1348	56.63	0.001 (-)	103	17.03	0.001 (-)	113	42.48	0.001 (-)	125	50.56	0.008 (-)	5123	52.36
Degraded humid forest	1367	19.83	0.001 (+)	379	15.92	0.001 (+)	242	39.83	0.001 (+)	120	45.03	0.001 (+)	95	38.67	0.001	1961	20.04
Total	6891	100		2382	100		606	100		265	100		246	100		9784	100

+, habitat is selected; -, habitat is avoided; significances are neglected when the given habitat had less than ten observed locations; x, habitat was not represented in any random point of the corresponding wintering event.

The pristine humid forests of Madagascar are multi-layered evergreen woodlands with a canopy that can reach 30–35 m high. Since the 1970s, 33.4% of this forest has been destroyed, and repeated clearing has often made regeneration impossible. Currently, 39% of the area of pristine forest is formally protected (Moat & Smith 2007). The structure and ecology of degraded humid forests varies enormously, ranging from nearly intact forest to wooded grassland open areas (Moat & Smith 2007).

Our results showed that Eleonora's Falcons selected degraded humid forests close to pristine humid forests. Although we cannot rule out the possibility that falcons hunt within pristine forests, our results suggest that they prefer forest edges, where they could perhaps be benefiting from a spill-over edge effect of prey insects. This could make these ecotonal areas more profitable for prey capture, with pristine humid forests acting as a source of prey (Rand *et al.* 2006) and the adjacent areas acting as sinks (Gibson *et al.* 2011).

In agreement with our findings, Bildstein (2006) stated that migratory raptors wintering in the neotropics prefer edges and second-growth forests, rather than large pristine ones (Morel & Morel 1992) and, even within Madagascar, Watson *et al.* (2004) observed a 'significantly greater number of raptors on the edge of the littoral forest than in the core'. It is likely, therefore, that the patches of pristine forests surrounded by a matrix of open areas could be exploited by Eleonora's Falcons, especially the edges between these two types of habitat. Finally, the results also showed a slight preference for cultivated areas, probably rice fields, where flying insects are abundant and easy to catch (Zefania 2001).

Conservation implications

It has been suggested that the loss of wintering habitat could negatively affect a population even more than the loss of breeding habitat (Sutherland 1996). Thus, conservation efforts addressed only to one season or stage of the life-cycle of this long-distance migratory species may be inefficient, being jeopardized by threats occurring in ranges completely different from the breeding ones (Martin *et al.* 2007).

Our results suggest that the primeval humid forests of Madagascar and the close-by ecotonal areas are the most important habitat for Eleonora's Falcons during the wintering season. The findings of the present study should be of concern in the light of the rapid decline of humid

forests in Madagascar, which could disappear by 2067 according to current predictions (Moat & Smith 2007). Our findings, although based on a limited sample, fill an important gap in the knowledge of the ecology of Eleonora's Falcons and could be used to focus conservation efforts and further field studies. In a global context, pristine forests currently attract the strongest protection efforts (Bermingham *et al.* 2005, Gibson *et al.* 2011), but our results, which are in agreement with other recent research on the value of tropical secondary forest (Chazdon *et al.* 2009), show that second growth forests, peripheral and ecotonal areas, especially when close to more pristine areas, deserve attention for the conservation of Eleonora's Falcons.

ACKNOWLEDGEMENTS

The Terra Natura Foundation and the 'Servicio de Biodiversidad' from the 'Conselleria de Medio Ambiente, Agua, Urbanismo y Vivienda' (Generalitat Valenciana) funded this project. Special thanks are due to all the people who helped in bird trapping. We also thank C. García-Ripollés, an anonymous referee and Phil Whitfield for their useful comments on an earlier draft of the manuscript. U. Mellone is supported by FPU grant of the Spanish Ministry of Education (reference AP2008-0947). R. Limiñana has a postdoctoral grant (reference 10/12-C) co-funded by 'Consejería de Educación y Ciencia' (JCCM) and the European Social Fund. P. López-López is supported by a 'Juan de la Cierva' postdoctoral grant of the Spanish Ministry of Science and Innovation (reference JCI-2011-09588). This paper is part of the PhD dissertation of U. Mellone at the University of Alicante. This study complies with the current laws in Spain.

REFERENCES

- Alerstam, T., Hake, M. & Kjellén, N.** 2006. Temporal and spatial patterns of repeated migratory journeys by ospreys. *Anim. Behav.* **71**: 555–566.
- Beyer, H.L.** 2004. Hawth's analysis tools for ArcGIS Available at: <http://www.spatial ecology.com/htools> (accessed 20 April 2010).
- Bermingham, E., Dick, C.W. & Moritz, C.** 2005. *Tropical Rain Forests: Past, Present and Future*. Chicago, IL: University of Chicago Press.
- Berthold, P.** 2001. *Bird Migration. A General Survey*. New York: Oxford University Press.
- Bildstein, K.L.** 2006. *Migrating Raptors of the World*. Ithaca, NY: Cornell University Press.
- BirdLife International** 1999. *International Species Action Plan: Eleonora's Falcon Falco eleonorae*.
- BirdLife International** 2010. Eleonora's falcon Species Action Plan implementation review. Available at: <http://www.birdlifeforums.org/WebX/.2cba720c> (accessed 20 October 2010).
- Bowlin, M.S., Bisson, I-A., Shamoun-Baranes, J., Reichard, J.D., Sapir, N., Marra, P.P., Kunz, T.H., Wilcove, D.S., Hedenström, A., Guglielmo, C.G., Åkesson, S.,**

- Ramenofsky, M. & Wikelski, M. 2010. Grand challenges in migration biology. *Integr. Comp. Biol.* **50**: 261–279.
- Cadahía, L., López-López, P., Urios, V. & José Negro, J.J. 2010. Satellite telemetry reveals individual variation in juvenile Bonelli's eagle dispersal areas. *Eur. J. Wildl. Res.* **56**: 923–930.
- Chazdon, R.L., Peres, C.A., Dent, D., Sheil, D., Lugo, A.E., Lamb, D., Stork, N.E. & Miller, S.E. 2009. The potential for species conservation in tropical secondary forests. *Conserv. Biol.* **23**: 1406–1417.
- Del Moral, J.C. 2008. *El Halcón de Eleonora en España. Población en 2004–2007 y método de censo*. Madrid: SEO/BirdLife.
- Dimalexis, A., Xirouchakis, S., Portolou, D., Latsoudis, P., Karris, G., Fric, J., Georgiakakis, P., Barboutis, C., Bourdakis, S., Ivovic, M., Kominos, T. & Kakalis, E. 2008. The status of Eleonora's Falcon (*Falco eleonorae*) in Greece. *J. Ornithol.* **149**: 23–30.
- García-Ripollés, C., López-López, P. & Urios, V. 2010. First description of migration and wintering of adult Egyptian vultures *Neophron percnopterus* tracked by GPS satellite telemetry. *Bird Study* **57**: 261–265.
- Gerkmann, B. & Meyburg, B.-U. 2009. Habitats used by Lesser Spotted Eagles (*Aquila pomarina*) during migration and wintering as revealed by Satellite tracking and remote sensing. *Popul. Ecol. Raptors. Owls.* **6**: 87–102.
- Gibson, L., Lee, T.M., Koh, L.P., Brook, B.W., Gardner, T.A., Barlow, J., Peres, Bradshaw C.J.A., Laurance, W.F., Lovejoy, T.E. & Sodhi, N.S. 2011. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**: 378–381.
- Gschweng, M., Kalko, E.K.V., Querner, U., Fiedler, W. & Berthold, P. 2008. All across Africa: highly individual migration routes of Eleonora's falcon. *Proc. R. Soc. Lond. B.* **275**: 2887–2896.
- Harper, G., Steinger, M., Tucker, C., Juhn, D. & Hawkins, F. 2007. Fifty years of deforestation and forest fragmentation in Madagascar. *Environ. Conserv.* **34**: 325–333.
- Hobson, K.A. 2008. Using endogenous and exogenous markers in bird conservation. *Bird Conserv. Int.* **18**: 174–199.
- Hood, G.M. 2010. *Poptools*, version 3.1. Available at: <http://www.cse.csiro.au/poptools> (accessed 10 October 2010).
- Hooge, P.N. & Eichenlaub, B. 2000. *Animal Movement Extension to Arcview*, version 2.0. Anchorage, AK: US Geological Survey.
- Irwin, M.T., Wright, P.C., Birkinshaw, C., Fisher, B., Gardner, C.J., Glos, J., Goodman, S.M., Loiselle, P., Rabeson, P., Raharison, J.L., Raterilalao, M.J., Rakotondravy, D., Raselimanana, A., Ratsimbazafy, J., Sparks, J., Wilmé, L. & Ganzhorn, J.U. 2010. Patterns of species change in anthropogenically disturbed forests of Madagascar. *Biol. Conserv.* **143**: 2351–2362.
- Jiguet, F., Barbet-Massin, M. & Chevallier, D. 2011. Predictive distribution models applied to satellite tracks: modelling the western African winter range of European migrant black storks. *J. Ornithol.* **152**: 111–118.
- Kassara, C., Fric, J. & Sfenthourakis, S. 2011. Home-range and resource use by Eleonora's falcon (*Falco eleonorae*) in its wintering quarters inferred by satellite telemetry data. In Fusani L., Coppack T. & Strazd M. (eds), *Proceedings of the 8th Conference of the European Ornithologists' Union*: 181. Latvian Ornithological Society, Riga, Latvia. Available at: <http://www.unife.it/dipartimento/biologia-evoluzione/progetti/eou2011/EOUabstractweb.pdf> (accessed 5 September 2011).
- Limiñana, R., Soutullo, A., López-López, P. & Urios, V. 2008. Pre-migratory movements of adult Montagu's Harriers *Circus pygargus*. *Ardea* **96**: 81–90.
- Limiñana, R., Soutullo, A., Urios, V. & Reig-Ferrer, A. 2011. Migration and wintering areas of adult Montagu's Harriers (*Circus pygargus*) breeding in Spain. *J. Ornithol.* online first: DOI: 10.1007/s10336-011-0698-x
- López-López, P., Limiñana, R. & Urios, V. 2009. Autumn migration of Eleonora's falcon *Falco eleonorae* tracked by satellite telemetry. *Zool. Stud.* **48**: 485–491.
- López-López, P., Limiñana, R., Mellone, U. & Urios, V. 2010. From the Mediterranean Sea to Madagascar. Are there ecological barriers for the long-distance migrant Eleonora's falcon? *Landsc. Ecol.* **25**: 803–813.
- Manly, B.F.J. 1997. *Randomization, Bootstrap and Monte Carlo Methods in Biology*, 2nd edn. London: Chapman & Hall.
- Martin, T.G., Chadès, I., Arcese, P., Marra, P.P., Possingham, H.P. & Ryan Norris, D. 2007. Optimal conservation of migratory species. *PLoS ONE* **2**(8): e751.
- Mellone, U., López-López, P., Limiñana, R. & Urios, V. 2011. Weather conditions promote route flexibility during open ocean crossing in a long-distance migratory raptor. *Int. J. Biometeorol.* **55**: 463–468.
- Mellone, U., Yáñez, B., Limiñana, R., Muñoz, A.R., Pavón, D., González, J.M., Urios, V. & Ferrer, M. 2011. Summer staging areas of non-breeding Short-toed Eagles. *Bird Study* iFirst: DOI: 10.1080/00063657.2011.598914
- Moat, J. & Smith, P. 2007. *Atlas of the Vegetation of Madagascar*. London: Kew Publishing.
- Morel, G.J. & Morel, M.Y. 1992. Habitat use by Palearctic migrant passerine birds in West Africa. *Ibis* **134**: 83–88.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., de Fonseca, G.A.B. & Kent, J. 2000. Biodiversity and hotspots for conservation priorities. *Nature* **403**: 853–858.
- Newton, I. 2004. Population limitation in migrants. *Ibis* **146**: 197–226.
- Rand, T.A., Tyliaakis, J.M. & Tschardtke, T. 2006. Spillover edge effects: the dispersal of agriculturally subsidized insect natural enemies into adjacent natural habitats. *Ecol. Lett.* **9**: 603–614.
- Rodríguez, A., Negro, J.J., Bustamante, J., Fox, J.W. & Afanasyev, V. 2009. Geolocators map the wintering grounds of threatened Lesser Kestrels in Africa. *Divers. Distrib.* **15**: 1010–1016.
- Sanderson, F.J., Donald, P.F., Pain, D.J., Burfield, I.J. & van Bommel, F.P.J. 2006. Long-term population declines in Afro-Palearctic migrant birds. *Biol. Conserv.* **131**: 93–105.
- Sergio, F., Newton, I. & Marchesi, L. 2005. Top predators and biodiversity. *Nature* **236**: 192.
- Sergio, F., Caro, T., Brown, D., Clucas, B., Hunter, J., Ketchum, J., McHugh, K. & Hiraldo, F. 2008. Top predators as conservation tools: ecological rationale, assumptions, and efficacy. *Annu. Rev. Ecol. Syst.* **39**: 1–19.
- Silverman, B.W. 1986. *Density Estimation for Statistics and Data Analysis*. Chapman and Hall, London.
- Soutullo, A., Cadahía, L., Urios, V., Ferrer, M. & Negro, J.J. 2007. Accuracy of lightweight satellite telemetry: a case study in the Iberian Peninsula. *J. Wildl. Manage.* **71**: 1010–1015.
- Soutullo, A., Urios, V., Ferrer, M. & López-López, P. 2008. Habitat use by juvenile Golden Eagles *Aquila chrysaetos* in Spain. *Bird Study* **55**: 236–240.
- Strandberg, R., Klaassen, R.H.G., Hake, M., Olofsson, P., Thorup, K. & Alerstam, T. 2008. Complex temporal pattern of Marsh Harrier *Circus aeruginosus* migration due to pre- and post-migratory movements. *Ardea* **96**: 159–171.
- Sutherland, W.J. 1996. Predicting the consequences of habitat loss for migratory populations. *Proc. R. Soc. Lond. B.* **263**: 1325–1327.
- Thorstrom, R. & René de Roland, L.-A. 2000. Status and conservation of raptors on the Masoala Peninsula, Madagascar. In Chancellor,

- R.D. & Meyburg, B.-U. (eds), *Raptors at Risk*: 35–41, World Working Group on Birds of Prey, Berlin; Hancock House, Blaine, WA.
- Walter, H.** 1979. *Eleonora's Falcon. Adaptations to Prey and Habitat in a Social Raptor*. London: The University of Chicago Press.
- Watson, J.E.M., Whittaker, R.J. & Dawson, T.P.** 2004. Habitat structure and proximity to forest edge affect the abundance and distribution of forest-dependent birds in tropical coastal forests of south-eastern Madagascar. *Biol. Conserv.* **120**: 311–327.
- Wikelski, M., Kays, R., Kasdin, J., Thorup, K., Smith, J.A., Cochran, W.W. & Swenson, G.W. Jr.** 2007. Going wild – what a global small-animal tracking system could do for experimental biologists. *J. Exp. Biol.* **210**: 181–186.
- Wolda, H.** 1978. Seasonal fluctuations in rainfall, food and abundance of tropical insects. *J. Anim. Ecol.* **47**: 369–381.
- Worton, B.J.** 1989. Kernel methods for estimating the utilization distribution in home-range studies. *Ecology* **70**: 164–168.
- Worton, B.J.** 1995. Using Monte Carlo simulation to evaluate kernel-based home range estimators. *J. Wildl. Manage.* **59**: 794–800.
- Zefania, S.** (ed.). 2001. Observation of sooty and Eleonora's falcons in Madagascar. In Anonymous, *Wings Over Africa. Proceedings of the International Seminar on Bird Migration: Research, Conservation, Education and Flight Safety*: 151–159, Society for Protection of Nature, Tel Aviv, Israel.

(MS received 24 June 2011; revised MS accepted 12 October 2011)